



Getting started with VEPPANU

As you begin your journey with VEPPANU, this guide can help you get the most benefit from your medication—understanding how this medication works, how to take it as prescribed, and what to expect once treatment begins.

Please see accompanying full Prescribing Information including Patient Information, or visit [VEPPANU.com](https://www.veppanu.com)

What is VEPPANU?

VEPPANU is a prescription medicine to treat people with estrogen receptor-positive, human epidermal growth factor receptor 2 (HER2)-negative, *estrogen receptor-1 (ESR1)*-mutated advanced breast cancer or breast cancer that has spread to other parts of the body (metastatic), **and** whose disease has progressed after at least one line of endocrine-based therapy.

Your healthcare provider will perform a test to make sure that VEPPANU is right for you.

It is not known if VEPPANU is safe and effective in children.

VEPPANU is the first and only FDA-approved PROTAC, or Proteolysis Targeting Chimera.

VEPPANU is an estrogen receptor (ER)-targeted hormone therapy that works directly with your body's machinery to degrade and remove ERs from cancer cells.

Please see accompanying full Prescribing Information including Patient Information, or visit [VEPPANU.com](https://www.VEPPANU.com)



What to do before taking VEPPANU

Tell your healthcare provider about all of your medical conditions, including if you:

- Have heart failure or heart rhythm problems, including QTc prolongation, and long QTc syndrome
- Have low blood levels of potassium or magnesium
- Are pregnant or plan to become pregnant. VEPPANU can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with VEPPANU
- Use effective birth control (contraception) during treatment with VEPPANU and for 2 weeks after the last dose
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with VEPPANU

Males with female partners who are able to become pregnant:

- Use effective birth control (contraception) during treatment with VEPPANU and for 2 weeks after the last dose
- Are breastfeeding or plan to breastfeed. It is not known if VEPPANU passes into your breast milk. Do not breastfeed during treatment with VEPPANU and for 2 weeks after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

VEPPANU and other medicines may affect the way each other works and may cause serious side effects. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

QTc=corrected QT interval

VEPPANU[™]
vepedegestrant 200 mg
100 mg

How should you take VEPPANU?

- Take VEPPANU exactly as your healthcare provider tells you
- Do not change your dose or stop taking VEPPANU unless your healthcare provider tells you



**Take VEPPANU
1 time each day**



**Take VEPPANU
with food**



**Swallow VEPPANU
tablets whole**

Do not chew, crush,
dissolve, or split the tablets
before swallowing



Do not take VEPPANU tablets that are broken,
cracked, or look damaged



If you miss a dose or vomit after taking
VEPPANU, do not take another dose of VEPPANU
on that day. Take your next dose the following
day at your regularly scheduled time. Do not
take 2 doses on the next treatment day

What should you avoid while taking VEPPANU?

Avoid taking St. John's wort, eating grapefruit, or drinking grapefruit juice during treatment with VEPPANU.

Please see accompanying full Prescribing
Information including Patient Information, or
visit VEPPANU.com

What are the possible side effects of VEPPANU?



VEPPANU can cause serious side effects, including:

- **Heart rhythm problems (QTc interval prolongation).**

VEPPANU can cause changes in the electrical activity of your heart and may increase your risk of abnormal heart rhythm problems, and sudden death. Your healthcare provider will check your heart with a test called an electrocardiogram (ECG) and check your blood potassium and magnesium levels before and as needed during treatment with VEPPANU. Get emergency medical help right away if you get any signs and symptoms of abnormal heart rhythm, including:

- Feeling lightheaded or faint
- Shortness of breath
- Dizziness
- Chest pain
- Feeling that your heart is pounding or beating fast (heart palpitations)

The most common side effects of VEPPANU include:

- Decreased white blood cell counts
- Increased liver function tests
- Muscle and bone pain
- Tiredness
- Decreased red blood cell counts
- Nausea
- Decreased potassium levels in your blood
- Decreased appetite
- Abnormal electrocardiogram (QT prolonged)
- Decreased platelet counts
- Constipation

VEPPANU
vepdegestrant 200 mg
100 mg

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with VEPPANU, if you develop certain side effects.

VEPPANU may affect fertility in males and in females who are able to become pregnant. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of VEPPANU.

Call your doctor for medical advice about side effects.
You may report side effects to FDA at 1-800-FDA-1088.

How should you store VEPPANU?



- Store VEPPANU at room temperature between 68°F to 77°F (20°C to 25°C)
- VEPPANU comes in a bottle with a child-resistant closure

Keep VEPPANU and all medicines out of the reach of children.

Please see accompanying full Prescribing Information including Patient Information, or visit VEPPANU.com

General information about the safe and effective use of VEPPANU



Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use VEPPANU for a condition for which it was not prescribed.

Do not give VEPPANU to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about VEPPANU that is written for health professionals.

RIGEL ONECARE can support you on your treatment journey with VEPPANU



The Patient Support Team is fully trained to support you with understanding your coverage, financial and copay assistance programs, and additional resources.



Scan the QR Code or visit RigelONECARE.com to learn how RIGEL ONECARE can support you

VEPPANU[™]
vepeggestrant 200 mg
100 mg



What to know about your treatment

- 1 VEPPANU is the first and only FDA-approved PROTAC, or **P**roteolysis **T**argeting **C**himeras. VEPPANU is for adult patients with ER+/HER2-, *ESR1*-mutated advanced or metastatic breast cancer, and whose disease has progressed after at least one line of endocrine-based therapy.
- 2 Take 1 VEPPANU tablet whole every day, by mouth, with food. Always take VEPPANU exactly as instructed.
- 3 Your healthcare provider will monitor your treatment for signs of heart rhythm problems (QTc interval prolongation).
- 4 There are other possible side effects, including decreased white blood cell counts, increased liver function tests, and muscle and bone pain.

Tell your healthcare provider about any health conditions and medications before starting treatment.



For more information visit:
VEPPANU.COM

Please see accompanying full Prescribing Information including Patient Information, or visit **VEPPANU.com**

© 2026 Rigel Pharmaceuticals, Inc.
All rights reserved. VEP_BC-26022 07/2026

VEPPANU is a trademark of Rigel Pharmaceuticals, Inc.
South San Francisco, CA 94080

